


Cytoplasmic Protein NCK1 (NCK1) Antibody

Catalogue No.:abx000952



Western blot analysis of lysates from mouse brain, using NCK1 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.

NCK1 Antibody is a Rabbit Polyclonal antibody against NCK1. The protein encoded by this gene is one of the signaling and transforming proteins containing Src homology 2 and 3 (SH2 and SH3) domains. It is located in the cytoplasm and is an adaptor protein involved in transducing signals from receptor tyrosine kinases to downstream signal recipients such as RAS. Alternatively spliced transcript variants encoding different isoforms have been found.

Target: Cytoplasmic Protein NCK1 (NCK1)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the

end user.

Conjugation: Unconjugated

Immunogen: Recombinant protein corresponding to NCK1. The exact sequence is proprietary.

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: P16333 (<u>UniProt</u>, <u>ExPASy</u>)

Gene Symbol: NCK1

Datasheet

Version: 4.0.0 Revision date: 12 Oct 2025



GeneID: <u>4690</u>

NCBI Accession: NP_006144.1

KEGG: hsa:4690

String: <u>9606.ENSP00000417273</u>

Molecular Weight: Calculated MW: 43 kDa

Observed MW: 43 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.