



## Bcl2 Associated X Protein Phospho-Thr167 (BAX pT167) Antibody

Catalogue No.:abx012466



BAX (Phospho-Thr167) antibody reacts with epitope-specific phosphopeptide and corresponding non-phosphopeptide. The absorbance readings at 450 nM are shown in the ELISA figure.



Immunofluorescence analysis of A549 cells, using BAX (Phospho-Thr167) antibody.

Rabbit polyclonal antibody against BAX protein. Immunogen region is C-terminal. Specificity is as follows for the reactive species: H:T167, M:T167, R:T167.

Target: Bcl2 Associated X Protein Phospho-Thr167 (BAX pT167)

Clonality: Polyclonal

Target Modification: Thr167

Modification: Phosphorylation

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, IF/ICC

Host: Rabbit

Recommended dilutions: IF/ICC: 1/100 - 1/500, ELISA: 1/40000. Optimal dilutions/concentrations should be determined by

the end user.

Conjugation: Unconjugated

**Immunogen:** The antiserum was produced against synthesized phosphopeptide derived from human Bax

around the phosphorylation site of threonine 167 (F-G-TP-P-T).

## **Datasheet**

Version: 2.0.0 Revision date: 18 Oct 2025



Isotype: IgG

Form: Liquid

**Purification:** Purified from rabbit antiserum by affinity chromatography using epitope-specific immunogen.

**Storage:** Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q07812 (UniProt, ExPASy)

Buffer: PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 0.02% sodium azide, 50% glycerol.

Concentration: 1 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.

2 of 2

Website: www.abbexa.com · Email: info@abbexa.com