



Bcl2-Associated Agonist Of Cell Death Phospho-Ser91/128 (BAD pS91/128) Antibody

Catalogue No.: abx012475



Western blot analysis of extracts from COS7 cells, treated with TNF- α (20ng/ml, 30mins), using BAD (Phospho-Ser91/128) antibody.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using BAD (Phospho-Ser91/128) antibody.

Rabbit polyclonal antibody against BAD protein. Immunogen region is Internal. Specificity is as follows for the reactive species: H:S91, M:S128, R:S129.

Target: Bcl2-Associated Agonist Of Cell Death Phospho-Ser91/128 (BAD pS91/128)

Clonality: Polyclonal

Target Modification: Ser91/128

Modification: Phosphorylation

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/3000, IHC: 1/50 - 1/100, ELISA: 1/10000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human BAD around the phosphorylation site of serine 91/serine 128 (E-L-SP-P-F).

Datasheet

Version: 2.0.0

Revision date: 15 Oct 2025



Isotype:	IgG
Form:	Liquid
Purification:	Purified from rabbit antiserum by affinity chromatography using epitope-specific immunogen.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q92934 (UniProt , ExPASy)
KEGG:	hsa:572
String:	9606.ENSP00000378040
Sequence:	CEEELSPFRG
Buffer:	PBS (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150 mM NaCl, 0.02% sodium azide, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only