



Nucleolar Transcription Factor 1 Phospho-Ser484 (UBTF pS484) Antibody

Catalogue No.: abx012712



Western blot analysis of extracts from HeLa cells, treated with calyculinA (50ng/ml, 30mins), using UBF (Phospho-Ser484) antibody.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue using UBF (Phospho-Ser484) antibody.

Rabbit polyclonal antibody against UBF protein. Immunogen region is Internal. Specificity is as follows for the reactive species: H:S484, M:S484, R:S484.

Target: Nucleolar Transcription Factor 1 Phospho-Ser484 (UBTF pS484)

Clonality: Polyclonal

Target Modification: Ser484

Modification: Phosphorylation

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/3000, IHC: 1/50 - 1/100, ELISA: 1/1000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: The antiserum was produced against synthesized phosphopeptide derived from human UBF around the phosphorylation site of serine 484 (P-E-SP-P-K).

Datasheet

Version: 4.0.0

Revision date: 20 Aug 2025



Isotype:	IgG
Form:	Liquid
Purification:	Purified from rabbit antiserum by affinity chromatography using epitope-specific immunogen.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P17480 (UniProt , ExPASy)
KEGG:	hsa:7343
String:	9606.ENSP00000302640
Sequence:	CRGKLPESPKRAE
Buffer:	PBS (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150 mM NaCl, 0.02% sodium azide, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only