



Proto-Oncogene C-Rel (REL) Antibody

Catalogue No.: abx001095



Western blot analysis of extracts of various cell lines using REL Antibody (1/1000 dilution).



Western blot analysis of extracts of various cell lines using REL Antibody (1/1000 dilution).

REL Antibody is a Rabbit Polyclonal antibody against REL. This gene encodes a protein that belongs to the Rel homology domain/immunoglobulin-like fold, plexin, transcription factor (RHD/IPT) family. Members of this family regulate genes involved in apoptosis, inflammation, the immune response, and oncogenic processes. This proto-oncogene plays a role in the survival and proliferation of B lymphocytes. Mutation or amplification of this gene is associated with B-cell lymphomas, including Hodgkin's lymphoma. Single nucleotide polymorphisms in this gene are associated with susceptibility to ulcerative colitis and rheumatoid arthritis. Alternative splicing results in multiple transcript variants encoding different isoforms.

Target: Proto-Oncogene C-Rel (REL)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: WB

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant fusion protein corresponding to human REL

Isotype: IgG

Datasheet

Version: 6.0.0
Revision date: 03 Oct 2025



Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q04864 (UniProt , ExPASy)
Gene Symbol:	REL
GeneID:	5966
NCBI Accession:	NP_002899.1
KEGG:	hsa:5966
String:	9606.ENSP00000295025
Molecular Weight:	Calculated MW: 65 kDa/68 kDa Observed MW: 78 kDa
Buffer:	PBS, pH 7.3, containing 0.01% thiomersal, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.