



Cyclic AMP-Dependent Transcription Factor ATF-1 (ATF1) Antibody

Catalogue No.: abx013416



Western blot analysis of extracts from COLO205 cells, HeLa cells and HepG2 cells, using ATF1 antibody.



Immunofluorescence analysis of HUVEC cells, using ATF1 antibody.

Rabbit polyclonal antibody against ATF1 protein. Immunogen region is C-terminal.

Target:	Cyclic AMP-Dependent Transcription Factor ATF-1 (ATF1)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 1/500 - 1/3000, IF/ICC: 1/100 - 1/500, ELISA: 1/5000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	The antiserum was produced against synthesized peptide derived from C-terminal of human ATF1.
Isotype:	IgG
Form:	Liquid
Purification:	Purified from rabbit antiserum by affinity chromatography using epitope-specific immunogen.

Datasheet

Version: 3.0.0

Revision date: 29 Sep 2025



Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P18846 (UniProt , ExPASy)
KEGG:	hsa:466
String:	9606.ENSP00000262053
Buffer:	PBS (without Mg^{2+} and Ca^{2+}), pH 7.4, 150 mM NaCl, 0.02% sodium azide, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only