



T-Cell-Specific Surface Glycoprotein CD28 (CD28) Antibody


Catalogue No.: abx129528




Western blot analysis of recombinant Mouse CD28 Protein.




Western blot analysis of Mouse Lymph node lysate, using Rabbit Anti-Mouse CD28 Antibody (5 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



IHC-P analysis of Mouse Cerebrum Tissue, with DAB staining, using Rabbit Anti-Mouse CD28 Antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).




IHC-P analysis of Mouse Spleen Tissue, with DAB staining, using Rabbit Anti-Mouse CD28 Antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis of Mouse Kidney Tissue, with DAB staining, using Rabbit Anti-Mouse CD28 Antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).

Datasheet

Version: 3.0.0
Revision date: 09 Apr 2025



IHC-P analysis of Mouse Testis Tissue, with DAB staining, using Rabbit Anti-Mouse CD28 Antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).

Cluster Of Differentiation 28 Antibody is a Rabbit Polyclonal against Cluster Of Differentiation 28.

Target:	T-Cell-Specific Surface Glycoprotein CD28 (CD28)
Clonality:	Polyclonal
Reactivity:	Mouse
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-5 µg/ml, IHC: 5-50 µg/ml, IF/ICC: 5-50 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx168377 - Recombinant CD28 (Asn20-Leu150) expressed in E. coli
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P31041 (UniProt , ExPASy)
KEGG:	mmu:12487
String:	10090.ENSMUSP00000027165
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.