



DNA Fragmentation Factor Subunit Alpha (DFFa) Antibody


Catalogue No.: abx129723




Western blot analysis of Rat Spleen lysate, using Rat SIT1 Antibody (2 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of the recombinant protein.




IHC-P analysis of Mouse Lung Tissue, with DAB staining.



IHC-P analysis of Mouse Lung Tissue, with DAB staining.



IHC-P analysis of Mouse Stomach Tissue, with DAB staining.



IHC-P analysis of Kidney Tissue, with DAB staining.

DNA Fragmentation Factor Subunit Alpha Antibody is a Rabbit Polyclonal against DNA Fragmentation Factor Subunit Alpha.

Target:	DNA Fragmentation Factor Subunit Alpha (DFFa)
Clonality:	Polyclonal
Reactivity:	Mouse
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	DFFa (Lys19-Ser262)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.