



Intercellular Adhesion Molecule 2 / CD102 (ICAM2) Antibody


Catalogue No.: abx130334




Western blot analysis of (1) Human Jurkat Cells and (2) Human HeLa cells.




Western blot analysis of recombinant Rat ICAM2.



IHC-P analysis of Rat Stomach Tissue, with DAB staining.




IHC-P analysis of Rat Intestine Tissue, with DAB staining.



IHC-P analysis of Rat Heart Tissue, with DAB staining.

Datasheet

Version: 3.0.0
Revision date: 14 Oct 2025



IHC-P analysis of Rat Liver Tissue, with DAB staining.

Intercellular Adhesion Molecule 2 Antibody is a Rabbit Polyclonal against Intercellular Adhesion Molecule 2.

Target:	Intercellular Adhesion Molecule 2 / CD102 (ICAM2)
Clonality:	Polyclonal
Reactivity:	Rat
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-5 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	ICAM2 (Val29-Phe247)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.