



Sodium- and Chloride-Dependent Taurine Transporter (SLC6A6) Antibody

Catalogue No.: abx014892



Western blot analysis of extracts from COLO205 cells, using SC6A6 antibody.



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using SC6A6 antibody.

Rabbit polyclonal antibody against SLC6A6 protein. Immunogen region is C-terminal.

Target: Sodium- and Chloride-Dependent Taurine Transporter (SLC6A6)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/3000, IHC: 1/50 - 1/100, ELISA: 1/10000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: The antiserum was produced against synthesized peptide derived from C-terminal of human SC6A6.

Isotype: IgG

Form: Liquid

Datasheet

Version: 2.0.0

Revision date: 05 Sep 2025



Purification: Purified from rabbit antiserum by affinity chromatography using epitope-specific immunogen.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: P31641 ([UniProt](#), [ExPASy](#))

KEGG: hsa:6533

String: [9606.ENSP00000480890](#)

Sequence: CREPNRWAVEREG

Buffer: PBS (without Mg^{2+} and Ca^{2+}), pH 7.4, 150 mM NaCl, 0.02% sodium azide, 50% glycerol.

Concentration: 1 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.