



Oncostatin-M (OSM) Antibody


Catalogue No.:abx128672



Western blot analysis of the recombinant protein.



IHC-P analysis of Human Stomach Tissue, with DAB staining.



IHC-P analysis of stomach tissue, with DAB staining.

Oncostatin-M Antibody is a Rabbit Polyclonal against Oncostatin-M.

Target: Oncostatin-M (OSM)

Research Area: Tumor Immunity

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Datasheet

Version: 4.0.0
Revision date: 20 Aug 2025



Conjugation:	Unconjugated
Immunogen:	OSM (Ala26-Arg220)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P13725 (UniProt , ExPASy)
Gene Symbol:	OSM
GeneID:	5008
KEGG:	hsa:5008
String:	9606.ENSP00000215781
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.