



Programmed Cell Death Protein 1 Ligand 2 (PDCD1LG2) Antibody


Catalogue No.: abx128134




Western blot analysis of recombinant Human PDCD1LG2.



Western blot analysis of Human Urine, using Rabbit Anti-Human PDCD1LG2 Antibody (2 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



IHC-P analysis of Human Tonsil Tissue, with DAB staining using Rabbit Anti-Human PDCD1LG2 Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis of Human Thyroid Cancer Tissue, with DAB staining using Rabbit Anti-Human PDCD1LG2 Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).

Programmed Cell Death Protein 1 Ligand 2 Antibody is a Rabbit Polyclonal against Programmed Cell Death Protein 1 Ligand 2.

Target: Programmed Cell Death Protein 1 Ligand 2 (PDCD1LG2)

Clonality: Polyclonal

Reactivity: Human

Datasheet

Version: 5.0.0
Revision date: 05 Mar 2025



Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx166054](#) - Recombinant PDCD1LG2 (Glu33-Phe236) expressed in E. coli

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q9BQ51 ([UniProt](#), [ExPASy](#))

String: [9606.ENSP00000380855](#)

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Concentration: 1 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.