



MAPK/MAK/MRK Overlapping Kinase (MOK) Antibody


Catalogue No.: abx129598




Western blot analysis of recombinant Mouse RAGE protein.




Western blot analysis of Human HepG2 cell lysate, using Mouse RAGE Antibody (3 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



IHC-P analysis of Mouse Liver tissue, with DAB staining, using Rabbit Anti-Mouse RAGE Antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis of Mouse Kidney tissue, with DAB staining, using Rabbit Anti-Mouse RAGE Antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis of Mouse Cardiac Muscle tissue, with DAB staining, using Rabbit Anti-Mouse RAGE Antibody (10 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).

Datasheet

Version: 3.0.0
Revision date: 29 Sep 2025



Renal Tumor Antigen Antibody is a Rabbit Polyclonal against Renal Tumor Antigen.

Target:	MAPK/MAK/MRK Overlapping Kinase (MOK)
Clonality:	Polyclonal
Reactivity:	Mouse
Tested Applications:	ELISA, WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	ELISA: 1/100 - 1/5000, WB: 1/50 - 1/400, IHC-P: 1/10 - 1/100, IHC-F: 1/50 - 1/500, IF/ICC: 1/10 - 1/100. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx168416 - Recombinant RAGE (Lys2-Lys228) expressed in E. coli
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Specificity:	Cross-reacts with Human and Rat RAGE.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.