



Signal Transducer And Activator of Transcription 2 (STAT2) Antibody


Catalogue No.:abx130350




Western blot analysis of (1) Rat Serum and (2) Mouse Lung Tissue.




Western blot analysis of recombinant Rat STAT2.



IHC-P analysis of Rat Adrenal Gland Tissue, with DAB staining.




IHC-P analysis of Rat Liver Tissue, with DAB staining.



IHC-P analysis of Rat Spinal Cord Tissue, with DAB staining.

Datasheet

Version: 2.0.0
Revision date: 13 Aug 2025



IHC-P analysis of Rat Stomach Tissue, with DAB staining.

STAT2 Antibody is a Rabbit Polyclonal against STAT2.

Target:	Signal Transducer And Activator of Transcription 2 (STAT2)
Clonality:	Polyclonal
Reactivity:	Rat
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	STAT2 (Arg587-Ile838)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q5XI26 (UniProt , ExPASy)
String:	10116.ENSARNOP00000046028
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.