



NAD-Dependent Protein Deacetylase Sirtuin-2 (SIRT2) Antibody


Catalogue No.: abx129627



WB analysis of Mouse brain (1), Mouse heart (2) and Rat brain tissue lysates (3) using SIRT2 antibody.



IHC-P analysis with DAB staining, of Mouse cerebrum tissue, using SIRT2 antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis with DAB staining, of Mouse cardiac muscle tissue, using SIRT2 antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).

NAD-Dependent Protein Deacetylase Sirtuin-2 (SIRT2) Antibody is a Rabbit Polyclonal antibody against NAD-Dependent Protein Deacetylase Sirtuin-2 (SIRT2).

Target: NAD-Dependent Protein Deacetylase Sirtuin-2 (SIRT2)

Research Area: Signal Transduction, Developmental Science

Clonality: Polyclonal

Reactivity: Mouse

Tested Applications: WB, IHC

Host: Rabbit

Datasheet

Version: 5.0.0
Revision date: 30 Aug 2025



Recommended dilutions: WB: 0.01-5 µg/ml, IHC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx167482](#) - Recombinant Mouse SIRT2 (Glu65-Glu340) expressed in E. coli.

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q8VDQ8 ([UniProt](#), [ExPASy](#))

Gene Symbol: SIRT2

GenelD: [64383](#)

KEGG: mmu:64383

String: [10090.ENSMUSP00000072732](#)

Buffer: PBS, pH 7.4, containing 0.02% NaN₃, 50% glycerol.

Specificity: Cross-reacts with Human and Rat SIRT2.

Concentration: 1 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.