



NAD-Dependent Protein Deacetylase Sirtuin-3, Mitochondrial (SIRT3) Antibody


Catalogue No.: abx129703




Western blot analysis of Mouse Heart Tissue.



Western blot analysis of recombinant Mouse SIRT3.



IHC-P analysis of Mouse Kidney Tissue, with DAB staining.



IHC-P analysis of Mouse Liver Tissue, with DAB staining.

NAD-Dependent Protein Deacetylase Sirtuin-3, Mitochondrial Antibody is a Rabbit Polyclonal antibody against NAD-Dependent Protein Deacetylase Sirtuin-3, Mitochondrial.

Target: NAD-Dependent Protein Deacetylase Sirtuin-3, Mitochondrial (SIRT3)

Research Area: Signal Transduction

Datasheet

Version: 3.0.0
Revision date: 08 Sep 2025



Clonality:	Polyclonal
Reactivity:	Mouse
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-5 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	SIRT3 (Gly5-Arg257)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q8R104 (UniProt , ExPASy)
Gene Symbol:	SIRT3
KEGG:	mmu:64384
String:	10090.ENSMUSP00000026559
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.