



NAD-Dependent Protein Deacetylase Sirtuin-5, Mitochondrial (SIRT5) Antibody

Catalogue No.: abx104979



Western blot analysis of the recombinant protein.



IHC-P analysis of adrenal tissue, with DAB staining.

NAD-Dependent Protein Deacetylase Sirtuin-5, Mitochondrial Antibody is a Rabbit Polyclonal antibody against NAD-Dependent Protein Deacetylase Sirtuin-5, Mitochondrial.

Target: NAD-Dependent Protein Deacetylase Sirtuin-5, Mitochondrial (SIRT5)

Research Area: Signal Transduction

Clonality: Polyclonal

Reactivity: Human

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: SIRT5 (Ser37-Ser310)

Form: Liquid

Datasheet

Version: 5.0.0

Revision date: 07 Sep 2025



Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q9NXA8 (UniProt , ExPASy)
Gene Symbol:	SIRT5
GeneID:	23408
OMIM:	604483
HGNC:	14933
Ensembl:	ENSG00000124523
String:	9606.ENSP00000476228
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.