



Transthyretin (TTR) Antibody


Catalogue No.: abx129948




Western blot analysis of Pig Heart lysate, using Pig TTR Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).




Western blot analysis of Pig Serum, using Rabbit Anti-Pig TTR Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of Pig Liver lysate, using Rabbit Anti-Pig TTR Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of Porcine Spleen lysate, using Rabbit Anti-Porcine TTR Antibody (1 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 0.2 µg/ml).



Western blot analysis of the recombinant protein.

Datasheet

Version: 2.0.0
Revision date: 11 Sep 2025



Transthyretin Antibody is a Rabbit Polyclonal against Transthyretin.

Target:	Transthyretin (TTR)
Clonality:	Polyclonal
Reactivity:	Pig
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	TTR (Gly21-Leu150)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P50390 (UniProt , ExPASy)
KEGG:	ssc:397419
String:	9823.ENSSSCP00000022931
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.