



Transitional Endoplasmic Reticulum ATPase (VCP) Antibody


Catalogue No.:abx128451




Western blot analysis of (1) Human Lung Tissue, (2) Human HeLa cells, (3) Porcine Brain Tissue and (4) Rat Brain Tissue.




Western blot analysis of recombinant Human VCP.



IHC-P analysis of Human Kidney Tissue, with DAB staining.



IHC-P analysis of Human Stomach Tissue, with DAB staining.




IHC-P analysis of Human Stomach Cancer Tissue, with DAB staining.

Datasheet

Version: 3.0.0

Revision date: 31 Aug 2025



IHC-P analysis of Human Liver Tissue, with DAB staining.

Transitional Endoplasmic Reticulum ATPase Antibody is a Rabbit Polyclonal against Transitional Endoplasmic Reticulum ATPase.

Target:	Transitional Endoplasmic Reticulum ATPase (VCP)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	VCP (Gly125-Ile371)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P55072 (UniProt , ExPASy)
KEGG:	hsa:7415
String:	9606.ENSP00000351777
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Datasheet

Version: 3.0.0

Revision date: 31 Aug 2025



Note:

THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only