

Protein-Tyrosine Kinase 2-Beta (PYK2) Antibody

Catalogue No.:abx016050



Western blot analysis using PYK2 antibody against Raji (1), PMA induced THP-1 (2), Jurkat (3) and Ramos (4) cell lysate.



Immunohistochemical analysis of paraffin-embedded human Tonsil tissues using PYK2 antibody.

PYK2: PTK2B protein tyrosine kinase 2 beta, also known as PTK2B, PKB, PTK, CAKB, FAK2, FRNK.Entrez Protein NP_004094.It is a cytoplasmic protein tyrosine kinase which is involved in calcium-induced regulation of ion channels and activation of the map kinase signaling pathway. The encoded protein may represent an important signaling intermediate between neuropeptide-activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. The encoded protein undergoes rapid tyrosine phosphorylation and activation in response to increases in the intracellular calcium concentration, nicotinic acetylcholine receptor activation, membrane depolarization, or protein kinase C activation. This protein has been shown to bind CRK-associated substrate, nephrocystin, GTPase regulator associated with FAK, and the SH2 domain of GRB2. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Four transcript variants encoding two different isoforms have been found for this gene.

Target:	Protein-Tyrosine Kinase 2-Beta (PYK2)
Clonality:	Monoclonal
Reactivity:	Human
Tested Applications:	ELISA, WB, IHC
Host:	Mouse
Recommended dilutions:	ELISA: 1/10000, WB: 1/500 - 1/2000, IHC: 1/200 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated

Datasheet Version: 4.0.0 Revision date: 21 Jul 2025



Immunogen:	Purified recombinant fragment of PYK2 (aa815-997) expressed in E. coli.
Isotype:	IgG _{2a}
Form:	Liquid
Purification:	Unpurified ascites.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q14289 (UniProt, ExPASy)
Gene Symbol:	РТК2В
GenelD:	2185
OMIM:	601212
HGNC:	9612
KEGG:	hsa:2185
Ensembl:	ENSG00000120899
String:	9606.ENSP00000380638
Molecular Weight:	115.8 kDa
Buffer:	Ascitic fluid containing 0.03% sodium azide.
Concentration:	Not determined.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.