



Bradykinin Receptor B2 (BDKRB2) Antibody


Catalogue No.: abx006762



Western blot analysis of various lysates using BDKRB2 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.



Immunofluorescence analysis of HeLa cells using BDKRB2 Antibody at dilution of 1/100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using BDKRB2 Antibody at dilution of 1/100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution. Blue: DAPI for nuclear staining.

BDKRB2 Antibody is a Rabbit Polyclonal Antibody against BDKRB2.

Target: Bradykinin Receptor B2 (BDKRB2)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IF/ICC

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/2000, IF/ICC: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 3.0.0
Revision date: 21 Jun 2025



Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 302-391 of human BDKRB2.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P30411 (UniProt , ExPASy)
Gene Symbol:	BDKRB2
GeneID:	624
OMIM:	113503
NCBI Accession:	NP_000614.1
HGNC:	1030
KEGG:	hsa:624
Ensembl:	ENSG00000168398
String:	9606.ENSP00000450482
Molecular Weight:	Calculated MW: 44 kDa Observed MW: 44 kDa
Buffer:	PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.
Concentration:	0.661 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.