



Butyrophilin-Like Protein 2 (BTNL2) Antibody

Catalogue No.: abx006640



Western blot analysis of lysates from wild type (WT) and 293T cells transfected with BTNL2 using BTNL2 Antibody at 1/3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded Rat kidney using BTNL2 Antibody at dilution of 1/400 (40x lens). High pressure antigen retrieval performed in 0.01 M Citrate buffer (pH 6.0) prior to IHC staining.

BTNL2 Antibody is a Rabbit Polyclonal Antibody against BTNL2.

Target:	Butyrophilin-Like Protein 2 (BTNL2)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB, IHC
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/1000 - 1/5000, IHC-P: 1/100 - 1/500. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 246-455 of human BTNL2.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.

Datasheet

Version: 3.0.0
Revision date: 09 Apr 2025



Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q9UIR0 ([UniProt](#), [ExPASy](#))

Gene Symbol: BTNL2

GeneID: [56244](#)

NCBI Accession: NP_062548.1

String: [9606.ENSP00000390613](#)

Molecular Weight: Calculated MW: 50 kDa
Observed MW: 65 kDa

Buffer: PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only