


Cluster of Differentiation 320 (CD320) Antibody

Catalogue No.: abx123610



Western blot analysis of lysates from U-251MG cells, using CD320 Antibody at 1/2500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 30s.

CD320 Antibody is a Rabbit Polyclonal Antibody against CD320.

Target:	Cluster of Differentiation 320 (CD320)
Clonality:	Polyclonal
Reactivity:	Human, Mouse
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/200 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant protein corresponding to CD320. The exact sequence is proprietary.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q9NPF0 (UniProt , ExPASy)
Gene Symbol:	CD320
GenelD:	51293

Datasheet

Version: 4.0.0

Revision date: 06 Oct 2025



NCBI Accession: NP_057663.1

Molecular Weight: Calculated MW: 29 kDa
Observed MW: 58-70 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only