


Cell Division Cycle 25C Phospho-Ser216 (CDC25C pS216) Antibody

Catalogue No.: abx123326



Western blot analysis of lysates from HT-29 cells, using Phospho-cdc25C-S216 Antibody at 1/400 dilution. HT-29 cells were treated by λ -PP mixed solution (1ul) at 30 °C for 30 minutes. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.

cdc25C (pS216) Antibody is a Rabbit Polyclonal Antibody against cdc25C (pS216).

Target:	Cell Division Cycle 25C Phospho-Ser216 (CDC25C pS216)
Clonality:	Polyclonal
Target Modification:	Ser216
Modification:	Phosphorylation
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB, IHC
Host:	Rabbit
Recommended dilutions:	ELISA: 1 μ g/ml, WB: 1/100 - 1/500, IHC-P: 1/50 - 1/200. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	A synthetic phosphorylated peptide around S216 of human cdc25C.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P30307 (UniProt , ExPASy)

Datasheet

Version: 5.0.0
Revision date: 02 Oct 2025



Gene Symbol: CDC25C

GeneID: [995](#)

NCBI Accession: NP_001781.2

KEGG: hsa:995

String: [9606.ENSP00000321656](#)

Molecular Weight: Calculated MW: 53 kDa
Observed MW: 60 kDa

Buffer: PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only