


Chemokine C-X-C-Motif Ligand 9 (CXCL9) Antibody

Catalogue No.: abx123481



Western blot analysis of various lysates using CXCL9 Antibody at 1/3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.

CXCL9 Antibody is a Rabbit Polyclonal Antibody against CXCL9.

Target:	Chemokine C-X-C-Motif Ligand 9 (CXCL9)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 50 to the C-terminus of human CXCL9.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q07325 (UniProt , ExPASy)
Gene Symbol:	CXCL9
GeneID:	4283

Datasheet

Version: 2.0.0

Revision date: 14 Apr 2025



NCBI Accession: NP_002407.1

Molecular Weight: Calculated MW: 14 kDa
Observed MW: 14 kDa

Buffer: PBS, pH 7.3, containing 0.09% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only