


Eukaryotic Translation Initiation Factor 3 Subunit F (EIF3F) Antibody

Catalogue No.: abx007170



Western blot analysis of lysates from HeLa cells, using EIF3F Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 60s.

Immunohistochemistry analysis of paraffin-embedded Human colon tissue using EIF3F Antibody at a dilution of 1/200 (40x lens). High pressure antigen retrieval was performed in 0.01 M citrate buffer (pH 6.0) prior to IHC staining.

EIF3F Antibody is a Rabbit Polyclonal Antibody against EIF3F.

Target:	Eukaryotic Translation Initiation Factor 3 Subunit F (EIF3F)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB, IHC, IF/ICC, IP
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/1000, IHC-P: 1/50 - 1/200, IF/ICC: 1/50 - 1/200, IP: 0.5 µg - 4 µg antibody per 200 µg - 400 µg extracts of whole cells. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 88-357 of human EIF3F.
Isotype:	IgG
Form:	Liquid

Datasheet

Version: 2.0.0

Revision date: 23 May 2025



Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	O00303 (UniProt , ExPASy)
Gene Symbol:	EIF3F
GeneID:	8665
NCBI Accession:	NP_003745.1
KEGG:	hsa:8665
String:	9606.ENSP00000431800
Molecular Weight:	Calculated MW: 38 kDa Observed MW: 38 kDa
Buffer:	PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.
Concentration:	> 0.2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.