


Interleukin 4 Induced Protein 1 (IL4I1) Antibody

Catalogue No.: abx006138



Western blot analysis of various lysates using IL4I1 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 10s.

Western blot analysis of lysates from THP-1 cells, using IL4I1 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 180s.

IL4I1 Antibody is a Rabbit Polyclonal Antibody against IL4I1.

Target: Interleukin 4 Induced Protein 1 (IL4I1)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant protein corresponding to IL4I1. The exact sequence is proprietary.

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Datasheet

Version: 2.0.0

Revision date: 05 Oct 2025



Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q96RQ9 ([UniProt](#), [ExPASy](#))

Gene Symbol: IL4I1

GeneID: [259307](#)

NCBI Accession: NP_690863.1

KEGG: hsa:259307

String: [9606.ENSP00000472474](#)

Molecular Weight: Calculated MW: 63 kDa
Observed MW: 75 kDa

Buffer: PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only