



Insulin-Induced Gene 2 Protein (INSIG2) Antibody


Catalogue No.: abx123558




Western blot analysis of extracts of Mouse kidney using INSIG2 Antibody (1/1000 dilution).



Western blot analysis of extracts of various cell lines using INSIG2 Antibody (1/1000 dilution).



Immunohistochemistry of paraffin-embedded mouse brain using INSIG2 Antibody (1/200 dilution, 40x lens). High pressure antigen retrieval was performed in 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded rat ovary using INSIG2 Antibody (1/200 dilution, 40x lens). High pressure antigen retrieval was performed in 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

INSIG2 Antibody is a Rabbit Polyclonal Antibody against INSIG2.

Target: Insulin-Induced Gene 2 Protein (INSIG2)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Datasheet

Version: 1.0.0
Revision date: 28 Oct 2025



Tested Applications: WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC-P: 1/50 - 1/200. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: A synthetic peptide corresponding to human INSIG2

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q9Y5U4 ([UniProt](#), [ExPASy](#))

GeneID: [51141](#)

NCBI Accession: NP_057217.2

Molecular Weight: Calculated MW: 24 kDa
Observed MW: 30 kDa

Buffer: PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

Concentration: 1 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.