



NADH Ubiquinone Oxidoreductase Subunit A7 (NDUFA7) Antibody

Catalogue No.: abx006153



Western blot analysis of various lysates using [KO Validated] NDUFA7 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 30s.



Western blot analysis of lysates from wild type (WT) and NDUFA7 knockout (KO) 293T cells, using [KO Validated] NDUFA7 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 5s.

NDUFA7 Antibody is a Rabbit Polyclonal Antibody against NDUFA7.

Target: NADH Ubiquinone Oxidoreductase Subunit A7 (NDUFA7)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-113 of human NDUFA7.

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Datasheet

Version: 2.0.0
Revision date: 01 Jul 2025



Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: O95182 ([UniProt](#), [ExPASy](#))

Gene Symbol: NDUFA7

GeneID: [4701](#)

NCBI Accession: NP_004992.2

KEGG: hsa:4701

String: [9606.ENSP00000301457](#)

Molecular Weight: Calculated MW: 13 kDa
Observed MW: 13 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.