


RAD51 Paralog D (Rad51D) Antibody

Catalogue No.: abx007016



Western blot analysis of lysates from Jurkat cells, using Rad51D Antibody at 1/500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 30s.

Rad51D Antibody is a Rabbit Polyclonal Antibody against Rad51D.

Target:	RAD51 Paralog D (Rad51D)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant protein corresponding to Rad51D. The exact sequence is proprietary.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	O75771 (UniProt , ExPASy)
Gene Symbol:	RAD51D
GeneID:	5892

Datasheet

Version: 5.0.0

Revision date: 29 Sep 2025



NCBI Accession: NP_002869.3

KEGG: hsa:5892

String: [9606.ENSP00000466834](#)

Molecular Weight: Calculated MW: 35 kDa
Observed MW: 35 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only