


Long-Chain Fatty Acid Transport Protein 6 (SLC27A6) Antibody

Catalogue No.: abx005736



Western blot analysis of various lysates using SLC27A6 Antibody at 1/1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution.
Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.

SLC27A6 Antibody is a Rabbit Polyclonal Antibody against SLC27A6.

Target:	Long-Chain Fatty Acid Transport Protein 6 (SLC27A6)
Clonality:	Polyclonal
Reactivity:	Human, Mouse
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 370-619 of human SLC27A6.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q9Y2P4 (UniProt , ExPASy)
Gene Symbol:	SLC27A6
GeneID:	28965

Datasheet

Version: 4.0.0

Revision date: 04 May 2025



NCBI Accession: NP_054750.1

String: [9606.ENSP00000262462](#)

Molecular Weight: Calculated MW: 70 kDa
Observed MW: 70 kDa

Buffer: PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only