


SAM Pointed Domain-Containing Ets Transcription Factor (SPDEF) Antibody

Catalogue No.: abx006389



Western blot analysis of lysates from HT-29 cells, using SPDEF Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.

SPDEF Antibody is a Rabbit Polyclonal Antibody against SPDEF.

Target:	SAM Pointed Domain-Containing Ets Transcription Factor (SPDEF)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-335 of human SPDEF.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	O95238 (UniProt , ExPASy)
Gene Symbol:	SPDEF

Datasheet

Version: 4.0.0
Revision date: 12 Oct 2025



GeneID: [25803](#)

OMIM: [608144](#)

NCBI Accession: NP_036523.1

HGNC: 17257

KEGG: hsa:25803

Ensembl: ENSG00000124664

String: [9606.ENSP00000363149](#)

Molecular Weight: Calculated MW: 38 kDa
Observed MW: 50 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.