



Succinate-CoA Ligase GDP-Forming Beta Subunit (SUCLG2) Antibody


Catalogue No.: abx124447




Western blot analysis of extracts of various cell lines using SUCLG2 Antibody (1/3000 dilution).




Immunohistochemistry of paraffin-embedded Human liver cancer using SUCLG2 Antibody (1/100 dilution, 40x lens).



Immunohistochemistry of paraffin-embedded Human lung cancer using SUCLG2 Antibody (1/100 dilution, 40x lens).



Immunofluorescence analysis of C6 cells using SUCLG2 Antibody (1/100 dilution, 40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using SUCLG2 Antibody (1/100 dilution, 40x lens). Blue: DAPI for nuclear staining.

Datasheet

Version: 5.0.0
Revision date: 26 Aug 2025



SUCLG2 Antibody is a Rabbit Polyclonal Antibody against SUCLG2.

Target:	Succinate-CoA Ligase GDP-Forming Beta Subunit (SUCLG2)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 1/500 - 1/2000, IHC-P: 1/50 - 1/200, IF/ICC: 1/50 - 1/100. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein corresponding to human SUCLG2
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q96I99 (UniProt , ExPASy)
Gene Symbol:	SUCLG2
GeneID:	8801
NCBI Accession:	NP_003839.2
Molecular Weight:	Calculated MW: 46 kDa/47 kDa Observed MW: 47 kDa
Buffer:	PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.