


TRK-Fused Gene Protein (TFG) Antibody

Catalogue No.: abx007059



Western blot analysis of various lysates using TRK fused gene (TFG) Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 30s.

TFG Antibody is a Rabbit Polyclonal Antibody against TFG.

Target: TRK-Fused Gene Protein (TFG)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-240 of human TRK fused gene (TFG).

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q92734 ([UniProt](#), [ExPASy](#))

Gene Symbol: TFG

GeneID: [10342](#)

Datasheet

Version: 4.0.0

Revision date: 18 Mar 2025



NCBI Accession: NP_006061.2

KEGG: hsa:10342

String: [9606.ENSP00000240851](https://string-db.org/cgi/show.do?query=9606.ENSP00000240851)

Molecular Weight: Calculated MW: 43 kDa
Observed MW: 55/43 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only