


Transient Receptor Potential Cation Channel Subfamily M Member 8 (TRPM8) Antibody

Catalogue No.: abx006769



Western blot analysis of various lysates using TRPM8 Antibody at 1/500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 60s.

TRPM8 Antibody is a Rabbit Polyclonal Antibody against TRPM8.

Target:	Transient Receptor Potential Cation Channel Subfamily M Member 8 (TRPM8)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 980-1104 of human TRPM8.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q7Z2W7 (UniProt , ExPASy)
Gene Symbol:	TRPM8

Datasheet

Version: 3.0.0

Revision date: 12 Sep 2025



GeneID: [79054](#)

NCBI Accession: NP_076985.4

KEGG: hsa:79054

String: [9606.ENSP00000323926](#)

Molecular Weight: Calculated MW: 128 kDa
Observed MW: 126 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only