


WD Repeat Domain Phosphoinositide-Interacting Protein 2 (WIP12) Antibody

Catalogue No.: abx007018



Western blot analysis of various lysates using WIP12 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 3s.

Western blot analysis of lysates from Rat testis, using WIP12 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 30s.

WIP12 Antibody is a Rabbit Polyclonal Antibody against WIP12.

Target: WD Repeat Domain Phosphoinositide-Interacting Protein 2 (WIP12)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant protein corresponding to WIP12. The exact sequence is proprietary.

Isotype: IgG

Form: Liquid

Datasheet

Version: 5.0.0

Revision date: 02 Sep 2025



Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q9Y4P8 (UniProt , ExPASy)
Gene Symbol:	WIP12
GeneID:	26100
NCBI Accession:	NP_057087.2
String:	9606.ENSP00000288828
Molecular Weight:	Calculated MW: 49 kDa Observed MW: 49 kDa
Buffer:	PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	> 0.2 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.