


Mannose Receptor C Type 2 (MRC2) Antibody

Catalogue No.: abx135802



Western blot analysis of various lysates using MRC2 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 10s.

MRC2 Antibody is a Rabbit Polyclonal against MRC2.

Target:	Mannose Receptor C Type 2 (MRC2)
Clonality:	Polyclonal
Reactivity:	Human, Mouse
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 120-300 of human MRC2.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q9UBG0 (UniProt , ExPASy)
Gene Symbol:	MRC2
GeneID:	9902

Datasheet

Version: 3.0.0

Revision date: 06 Oct 2025



NCBI Accession: NP_006030.2

String: [9606.ENSP00000307513](#)

Molecular Weight: Calculated MW: 167 kDa
Observed MW: 190 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only