



Neprilysin / NEP (MME) Antibody (FITC)

Catalogue No.:abx139752



Surface staining of human peripheral blood cells with anti-CD10 FITC.



Fluorescence emission spectra of FITC.

CD10 Antibody is a Mouse Monoclonal against CD10.

Target: Neprilysin / NEP (MME)

Clonality: Monoclonal

Clone: C207

Reactivity: Human

Tested Applications: FCM

Host: Mouse

Recommended dilutions: FCM: 20 µl/100 µl of whole blood or 10⁶ cells. Optimal dilutions/concentrations should be determined by the end user. The content of a vial (2 ml) is sufficient for 100 tests.

Conjugation: FITC

Excitation/Emission: 499/515

Laser Line: 488

Immunogen: Mouse NALM-6 leukemia pre-B cell line (tissue/cell preparation).

Datasheet

Version: 1.0.0

Revision date: 10 Nov 2025



Isotype:	IgG ₁
Storage:	Store in the dark at 2-8°C. Avoid exposure to light. Do not freeze.
UniProt Primary AC:	P08473 (UniProt , ExPASy)
Gene Symbol:	MME
GeneID:	4311
KEGG:	hsa:4311
String:	9606.ENSP00000418525
Buffer:	Stabilizing PBS solution containing 15 mM sodium azide.
Specificity:	The antibody reacts with CD10 antigen (CALLA - Common acute lymphatic leukemia antigen), a 100 kDa type II integral membrane protein.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only