


DNA Damage-Binding Protein 2 (DDB2) Antibody

Catalogue No.:abx001528



Western blot analysis of various lysates using DDB2 Antibody at 1/500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.

DDB2 Antibody is a Rabbit Polyclonal antibody against DDB2. This gene encodes a protein that is necessary for the repair of ultraviolet light-damaged DNA. This protein is the smaller subunit of a heterodimeric protein complex that participates in nucleotide excision repair, and this complex mediates the ubiquitylation of histones H3 and H4, which facilitates the cellular response to DNA damage. This subunit appears to be required for DNA binding. Mutations in this gene cause xeroderma pigmentosum complementation group E, a recessive disease that is characterized by an increased sensitivity to UV light and a high predisposition for skin cancer development, in some cases accompanied by neurological abnormalities. Two transcript variants encoding different isoforms have been found for this gene.

Target: DNA Damage-Binding Protein 2 (DDB2)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB, IF/ICC

Host: Rabbit

Recommended dilutions: ELISA: 1 μg/ml, WB: 1/500 - 1/2000, IF/ICC: 1/50 - 1/200. Optimal dilutions/concentrations should

be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant protein corresponding to DDB2. The exact sequence is proprietary.

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

1 of 2

Website: www.abbexa.com · Email: info@abbexa.com

Datasheet

Version: 6.0.0 Revision date: 23 Aug 2025



UniProt Primary AC: Q92466 (UniProt, ExPASy)

Gene Symbol: DDB2

GeneID: <u>1643</u>

NCBI Accession: NP_000098.1

KEGG: hsa:1643

String: <u>9606.ENSP00000256996</u>

Molecular Weight: Calculated MW: 48 kDa

Observed MW: 48 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.

Website: www.abbexa.com · Email: info@abbexa.com