


Estradiol 17-Beta-Dehydrogenase 2 (HSD17B2) Antibody

Catalogue No.: abx001615



Western blot analysis of various lysates, using HSD17B2 Antibody at 1/500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 0.1s.

HSD17B2 Antibody is a Rabbit Polyclonal antibody against HSD17B2.

Target:	Estradiol 17-Beta-Dehydrogenase 2 (HSD17B2)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 100-200 of human HSD17B2.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P37059 (UniProt , ExPASy)
Gene Symbol:	HSD17B2
GeneID:	3294

Datasheet

Version: 4.0.0

Revision date: 11 Apr 2025



NCBI Accession: NP_002144.1

KEGG: hsa:3294

String: [9606.ENSP00000199936](#)

Molecular Weight: Calculated MW: 43 kDa
Observed MW: 35-40 kDa

Buffer: PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only