



Nuclear Factor, Erythroid Derived 2 Like 2 (NFE2L2) Antibody


Catalogue No.:abx126236




Western blot analysis of various lysates using NRF2 Antibody at ARC50393-02 dilution.HeLa cells were treated by MG132(50 µM) at 37 □ for 90 minutes.




Western blot analysis of various lysates using NRF2 Antibody at ARC50393-02 dilution.NIH3T3 cells were treated by MG132(50 µM) at 37 □ for 90 minutes.



Western blot analysis of various lysates using NRF2 antibody at dilution.C6 cells were treated by MG132(50 μ M) at 37 \square for 90 minutes.



Immunohistochemistry of paraffin-embedded human breast cancer using NRF2 antibody (1/100 dilution, 40x lens). High pressure antigen retrieval was performed in 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded human placenta using NRF2 antibody (1/100 dilution, 40x lens). High pressure antigen retrieval was performed in 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Datasheet

Version: 4.0.0 Revision date: 20 Aug 2025



NFE2L2 Antibody is a Rabbit Polyclonal against NFE2L2.

Target: Nuclear Factor, Erythroid Derived 2 Like 2 (NFE2L2)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC-P: 1/50 - 1/200. Not tested in IHC-F. Optimal dilutions/concentrations

should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant fusion protein corresponding to human NRF2

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: Q16236 (UniProt, ExPASy)

Gene Symbol: NFE2L2

GeneID: <u>4780</u>

NCBI Accession: NP_006155.2

KEGG: hsa:4780

String: <u>9606.ENSP00000380252</u>

Molecular Weight: Calculated MW: 65 kDa/66 kDa/67 kDa

Observed MW: 97-100 kDa

Buffer: PBS, pH 7.3, containing 0.05% BSA, 0.05% Proclin-300, 50% glycerol.

Concentration: 1 mg/ml

2 of 3

Datasheet

Version: 4.0.0 Revision date: 20 Aug 2025



Note:

THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.



3 of 3