


Solute Carrier Family 2, Facilitated Glucose Transporter Member 1 (SLC2A1) Antibody

Catalogue No.: abx126576



Western blot analysis of lysates from HepG2 cells, using GLUT1/SLC2A1 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 3min.

SLC2A1 Antibody is a Rabbit Polyclonal against SLC2A1.

Target:	Solute Carrier Family 2, Facilitated Glucose Transporter Member 1 (SLC2A1)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Synthetic peptide corresponding to SLC2A1. The exact sequence is proprietary.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P11166 (UniProt , ExPASy)
Gene Symbol:	SLC2A1

Datasheet

Version: 5.0.0

Revision date: 02 Oct 2025



GeneID: [6513](#)

NCBI Accession: NP_006507.2

KEGG: hsa:6513

String: [9606.ENSP00000416293](#)

Molecular Weight: Calculated MW: 54 kDa
Observed MW: 50 kDa

Buffer: PBS, pH 7.3, containing 0.01% thimerosal, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only