



Transforming Growth Factor Beta Receptor 2 (TGFB2) Antibody


Catalogue No.: abx126697




Western blot analysis of lysates from Mouse liver, using TGF beta Receptor II (TGFB2) Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.




Western blot analysis of lysates from Mouse liver using TGF beta Receptor II (TGFB2) Antibody at 1/1000 dilution incubated overnight at 4 °C. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded Human appendix using TGF beta Receptor II (TGFB2) Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer using TGF beta Receptor II (TGFB2) Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human breast using TGF beta Receptor II (TGFB2) Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.

Datasheet

Version: 4.0.0
Revision date: 06 Sep 2025



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using TGF beta Receptor II (TGFB2) Antibody at dilution of 1/100 (40x lens). Microwave antigen retrieval performed in 0.01 M PBS Buffer (pH 7.2) prior to IHC staining.

TGFB2 Antibody is a Rabbit Polyclonal against TGFB2.

Target:	Transforming Growth Factor Beta Receptor 2 (TGFB2)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/1000, IHC-P: 1/50 - 1/200, IF/ICC: 1/50 - 1/200. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant protein corresponding to TGFB2. The exact sequence is proprietary.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P37173 (UniProt , ExPASy)
Gene Symbol:	TGFB2
GeneID:	7048
NCBI Accession:	NP_003233.4

Datasheet

Version: 4.0.0

Revision date: 06 Sep 2025



KEGG: hsa:7048

String: [9606.ENSP00000351905](#)

Molecular Weight: Calculated MW: 65 kDa
Observed MW: 90 kDa

Buffer: PBS, pH 7.3, containing 0.01% thimerosal, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only