



ADAM Metallopeptidase Domain 10 (ADAM10) Antibody

Catalogue No.: abx125489



Western blot analysis of lysates from RAW264.7 cells, using ADAM10 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 20s.



Immunofluorescence analysis of HeLa cells using ADAM10 Antibody at dilution of 1/100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1/500 dilution. Blue: DAPI for nuclear staining.

ADAM10 Antibody is a Rabbit Polyclonal against ADAM10.

Target: ADAM Metallopeptidase Domain 10 (ADAM10)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB, IF/ICC

Host: Rabbit

Recommended dilutions: ELISA: 1 µg/ml, WB: 1/500 - 1/1000, IF/ICC: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant protein corresponding to ADAM10. The exact sequence is proprietary.

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Datasheet

Version: 4.0.0
Revision date: 28 Jul 2025



Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: O14672 ([UniProt](#), [ExPASy](#))

Gene Symbol: ADAM10

GeneID: [102](#)

NCBI Accession: NP_001101.1

KEGG: hsa:102

String: [9606.ENSP00000260408](#)

Molecular Weight: Calculated MW: 84 kDa
Observed MW: 90 kDa

Buffer: PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.