



Apoptosis Inducing Factor, Mitochondria Associated 2 (AIFM2) Antibody


Catalogue No.: abx125507




Western blot analysis of extracts of HepG2 cells using AIFM2/AMID/AMID/AMID Antibody (1/1000 dilution).



Western blot analysis of extracts of various cell lines using AIFM2/AMID/AMID/AMID Antibody (1/1000 dilution).



Immunohistochemistry of paraffin-embedded human breast cancer using AIFM2/AMID/AMID/AMID Antibody (1/20 dilution, 40x lens). High pressure antigen retrieval was performed in 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded human liver cancer using AIFM2/AMID/AMID/AMID Antibody (1/20 dilution, 40x lens). High pressure antigen retrieval was performed in 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

AIFM2 Antibody is a Rabbit Polyclonal against AIFM2.

Target: Apoptosis Inducing Factor, Mitochondria Associated 2 (AIFM2)

Clonality: Polyclonal

Datasheet

Version: 2.0.0
Revision date: 01 Jul 2025



Reactivity:	Human, Mouse, Rat
Tested Applications:	WB, IHC
Host:	Rabbit
Recommended dilutions:	WB: 1/500 - 1/2000, IHC-P: 1/50 - 1/200. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein corresponding to human AIFM2/AMID/AMID/AMID
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q9BRQ8 (UniProt , ExPASy)
Gene Symbol:	AIFM2
GeneID:	84883
NCBI Accession:	NP_116186.1
String:	9606.ENSP00000478931
Molecular Weight:	Calculated MW: 36 kDa/40 kDa Observed MW: 41 kDa
Buffer:	PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	1 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.