


## Butyrophilin Subfamily 2 Member A1 (BTN2A1) Antibody

Catalogue No.: abx125581



Western blot analysis of lysates from RD cells, using BTN2A1 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 180s.

BTN2A1 Antibody is a Rabbit Polyclonal against BTN2A1.

|                               |                                                                                                            |
|-------------------------------|------------------------------------------------------------------------------------------------------------|
| <b>Target:</b>                | Butyrophilin Subfamily 2 Member A1 (BTN2A1)                                                                |
| <b>Clonality:</b>             | Polyclonal                                                                                                 |
| <b>Reactivity:</b>            | Human                                                                                                      |
| <b>Tested Applications:</b>   | ELISA, WB                                                                                                  |
| <b>Host:</b>                  | Rabbit                                                                                                     |
| <b>Recommended dilutions:</b> | ELISA: 1 µg/ml, WB: 1/500 - 1/1000. Optimal dilutions/concentrations should be determined by the end user. |
| <b>Conjugation:</b>           | Unconjugated                                                                                               |
| <b>Immunogen:</b>             | Recombinant protein corresponding to BTN2A1. The exact sequence is proprietary.                            |
| <b>Isotype:</b>               | IgG                                                                                                        |
| <b>Form:</b>                  | Liquid                                                                                                     |
| <b>Purification:</b>          | Purified by affinity chromatography.                                                                       |
| <b>Storage:</b>               | Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.                                             |
| <b>UniProt Primary AC:</b>    | Q7KYR7 ( <a href="#">UniProt</a> , <a href="#">ExPASy</a> )                                                |
| <b>Gene Symbol:</b>           | BTN2A1                                                                                                     |
| <b>GeneID:</b>                | <a href="#">11120</a>                                                                                      |

# Datasheet

Version: 4.0.0

Revision date: 11 Oct 2025



**NCBI Accession:** NP\_510961.1

**KEGG:** hsa:11120

**String:** [9606.ENSP00000312158](#)

**Molecular Weight:** Calculated MW: 60 kDa  
Observed MW: 69 kDa

**Buffer:** PBS, pH 7.3, containing 0.05% Proclin-300, 50% glycerol.

**Concentration:** > 0.2 mg/ml

**Note:** THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only