


SLC16A10 Antibody

Catalogue No.: abx126560



Western blot analysis of various lysates using SLC16A10 Antibody at 1/1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution.
Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 10s.

SLC16A10 Antibody is a Rabbit Polyclonal against SLC16A10.

Target:	SLC16A10
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 456-515 of human SLC16A10.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q8TF71 (UniProt , ExPASy)
Gene Symbol:	SLC16A10
GeneID:	117247

Datasheet

Version: 3.0.0

Revision date: 31 Aug 2025



NCBI Accession: NP_061063.2

KEGG: hsa:117247

String: [9606.ENSPP00000357844](#)

Molecular Weight: Calculated MW: 55 kDa
Observed MW: 55 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only