



Activated Leukocyte Cell Adhesion Molecule / CD166 (ALCAM) Antibody

Catalogue No.: abx302383



IHC-P analysis of human liver tissue, using ALCAM antibody (1/100 dilution).



IF analysis of HepG2 cells, using ALCAM antibody (1/100 dilution) and AF488-conjugated Goat anti-Rabbit IgG (H+L).

ALCAM Antibody is a Rabbit Polyclonal against ALCAM.

Target: Activated Leukocyte Cell Adhesion Molecule / CD166 (ALCAM)

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: IHC: 1/20 - 1/200, IF/ICC: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant Human CD166 antigen protein (28-516 AA).

Isotype: IgG

Form: Liquid

Datasheet

Version: 7.0.0
Revision date: 15 Aug 2025



Purity:	> 95%
Purification:	Purified by Protein G.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q13740 (UniProt , ExPASy)
Gene Symbol:	ALCAM
GeneID:	214
OMIM:	601662
NCBI Accession:	NP_001230209.1, NM_001243280.1, NP_001230212.1, NM_001243283.1, NP_001618.2, NM_001627.3
KEGG:	hsa:214
String:	9606.ENSP00000305988
Buffer:	0.01 M PBS, pH 7.4, 0.03% Proclin-300 and 50% Glycerol.
Concentration:	5 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.