


Protein Max (MAX) Antibody

Catalogue No.:abx001769



Western blot analysis of lysates from mouse kidney, using MAX Antibody at 1/500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Exposure time: 150s.

MAX Antibody is a Rabbit Polyclonal antibody against MAX. The protein encoded by this gene is a member of the basic helix-loop-helix leucine zipper (bHLHZ) family of transcription factors. It is able to form homodimers and heterodimers with other family members, which include Mad, Mxi1 and Myc. Myc is an oncoprotein implicated in cell proliferation, differentiation and apoptosis. The homodimers and heterodimers compete for a common DNA target site (the E box) and rearrangement among these dimer forms provides a complex system of transcriptional regulation. Mutations of this gene have been reported to be associated with hereditary pheochromocytoma. A pseudogene of this gene is located on the long arm of chromosome 7. Alternative splicing results in multiple transcript variants.

Target: Protein Max (MAX)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat

Tested Applications: ELISA, WB

Host: Rabbit

 $\textbf{Recommended dilutions:} \ \ \textbf{ELISA: 1} \ \ \mu\text{g/ml, WB: 1/500 - 1/2000.} \ \ \text{Optimal dilutions/concentrations should be determined by the light of t$

end user.

Conjugation: Unconjugated

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-160 of human

MAX.

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

1 of 2

Datasheet

Version: 4.0.0 Revision date: 05 Mar 2025



UniProt Primary AC: P61244 (UniProt, ExPASy)

Gene Symbol: MAX

GeneID: <u>4149</u>

NCBI Accession: NP_002373.3

KEGG: hsa:4149

String: <u>9606.ENSP00000351490</u>

Molecular Weight: Calculated MW: 18 kDa

Observed MW: 19 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC,

THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL

CONSUMPTION.

2 of 2