



Sodium- and Chloride-Dependent Taurine Transporter (SLC6A6) Antibody

Catalogue No.: abx302899



WB analysis of (1) PC-3 whole cell lysates, (2) HL60 whole cell lysates, (3) mouse brain tissue, and (4) mouse liver tissue, using SLC6A6 antibody (1/1000 dilution). Predicted band size: 70 kDa, Observed band size: 70 kDa.



IF analysis of MCF7 cells, using SLC6A6 antibody (1/25 dilution) and AF488-conjugated Goat anti-Rabbit IgG (H+L).

SLC6A6 Antibody is a Rabbit Polyclonal against SLC6A6.

Target: Sodium- and Chloride-Dependent Taurine Transporter (SLC6A6)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IF/ICC: 1/20 - 1/200. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant human Sodium- and chloride-dependent taurine transporter protein (1-49 AA).

Isotype: IgG

Form: Liquid

Datasheet

Version: 4.0.0
Revision date: 05 May 2025



Purity:	> 95%
Purification:	Purified by Protein G.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P31641 (UniProt , ExPASy)
Gene Symbol:	SLC6A6
GeneID:	6533
OMIM:	186854
NCBI Accession:	NP_001127839.2, NM_001134367.3, NP_001127840.1, NM_001134368.3
KEGG:	hsa:6533
String:	9606.ENSP00000480890
Buffer:	0.01 M PBS, pH 7.4, 0.03% Proclin-300 and 50% Glycerol.
Concentration:	3.24 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.